

## PAPILIO XUTHUS.



### CHARACTER GENERICUS.

*Antennæ* apicem versus crassiores, sæpius clavato-capitatae.

*Alæ* (sedentis) erectæ sursumque conniventes, (volatu diurno.)

*Lin. Syst. Nat.* p. 774.

### CHARACTER SPECIFICUS, &c.

PAPILIO alis nigris albo striato maculatis; posterioribus subtus ocellis cærulcis fulvisque subfasciatis.

*Fab. sp. ins.* 2. p. 19.

PAPILIO alis caudatis nigris albido striato-maculatis; posticis subtus cæruleo fulvoque subocellatis.

*Lin. Syst. Nat.* p. 751.

---

Inter Lepidoptera rariora numeratur Papilio Xuthus, Indiæ orientalis incola. Tabula magnitudine vera pictum exhibet.







184

RN

# XUTHUS.



## GENERIC CHARACTER.

*Antennæ* thickening towards the upper part, and generally terminating in a clavated tip.

*Wings* (when sitting) erect, and meeting upwards. (Flight diurnal.)

## SPECIFIC CHARACTER, &c.

Black subcaudated Butterfly, with yellowish white striated spots; the lower wings marked beneath with a band of blue and orange ocellated spots.

---

The *Papilio Xuthus* is numbered among the rarer species of *Lepidoptera*, and is a native of the East Indies. The plate represents it in its natural size.